

Immunotag™ PLS2 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN1057
Product Description	Immunotag™ PLS2 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PLS2
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	Synthesized peptide derived from human protein . at AA range: 90-170
Specificity	PLS2 Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	PLSCR2
Accession No.	Q9NRY7 Q9DCW2

Antibody Specification

Description	phospholipid scramblase 2(PLSCR2) Homo sapiens This gene encodes a member of the phospholipid scramblase family. Phospholipid scramblases are membrane proteins that mediate calcium-dependent, non-specific movement of plasma membrane phospholipids and phosphatidylserine exposure. The encoded protein contains a low affinity calcium binding motif and may play a role in blood coagulation and apoptosis. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Jan 2011],
Protein Expression	Testis,
Subcellular Localization	nucleus,plasma membrane,integral component of membrane,
Protein Function	cofactor:Calcium.,function:May mediate accelerated ATP-independent bidirectional transbilayer migration of phospholipids upon binding calcium ions that results in a loss of phospholipid asymmetry in the plasma membrane. May play a central role in the initiation of fibrin clot formation, in the activation of mast cells and in the recognition of apoptotic and injured cells by the reticuloendothelial system.,online information:Scramblase entry,similarity:Belongs to the phospholipid scramblase family.,tissue specificity:Exclusively expressed in testis.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.