

# Immunotag™ RPF6 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN1149
Product Description	Immunotag™ RPF6 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	RPF6
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	Synthesized peptide derived from human protein . at AA range: 460-540
Specificity	RPF6 Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	RAPGEF6 PDZGEF2
Accession No.	Q8TEU7

## Antibody Specification

Description	function:Guanine nucleotide exchange factor (GEF) for Rap1A, Rap2A and M-Ras GTPases. Does not interact with cAMP.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 cyclic nucleotide-binding domain.,similarity:Contains 1 N-terminal Ras-GEF domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 Ras-associating domain.,similarity:Contains 1 Ras-GEF domain.,subcellular location:Upon binding to M-Ras, it translocates to the plasma membrane.,subunit:Interacts with the second PDZ domain of human PTP1e.,tissue specificity:Isoform 3 has highest expression levels in the brain, heart, liver, lung and placenta and is barely detectable in skeletal muscle, kidney and pancreas.,
Protein Expression	Brain,Clones donated by RIKEN,Epithelium,Fetal brain,PCR re
Subcellular Localization	intracellular,cytosol,plasma membrane,apical plasma membrane,endocytic vesicle,
Protein Function	function:Guanine nucleotide exchange factor (GEF) for Rap1A, Rap2A and M-Ras GTPases. Does not interact with cAMP.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 cyclic nucleotide-binding domain.,similarity:Contains 1 N-terminal Ras-GEF domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 Ras-associating domain.,similarity:Contains 1 Ras-GEF domain.,subcellular location:Upon binding to M-Ras, it translocates to the plasma membrane.,subunit:Interacts with the second PDZ domain of human PTP1e.,tissue specificity:Isoform 3 has highest expression levels in the brain, heart, liver, lung and placenta and is barely detectable in skeletal muscle, kidney and pancreas.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.