

Immunotag™ SPDEF Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN1246
Product Description	Immunotag™ SPDEF Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	SPDEF
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from human protein . at AA range: 140-220
Specificity	SPDEF Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	SPDEF PDEF PSE
Accession No.	O95238 Q9WTP3

Antibody Specification

Description	SAM pointed domain containing ETS transcription factor(SPDEF) Homo sapiens The protein encoded by this gene belongs to the ETS family of transcription factors. It is highly expressed in the prostate epithelial cells, and functions as an androgen-independent transactivator of prostate-specific antigen (PSA) promoter. Higher expression of this protein has also been reported in brain, breast, lung and ovarian tumors, compared to the corresponding normal tissues, and it shows better tumor-association than other cancer-associated molecules, making it a more suitable target for developing specific cancer therapies. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2011],
Protein Expression	Colon,Prostate,Prostatic carcinoma,
Subcellular Localization	nucleus,
Protein Function	function:May function as an androgen-independent transactivator of the prostate-specific antigen (PSA) promoter. Binds to 5'-GGAT-3' DNA sequences. May play a role in the regulation of the prostate gland and/or prostate cancer development. Acts as a transcriptional activator for SERPINB5 promoter.,similarity:Belongs to the ETS family.,similarity:Contains 1 ETS DNA-binding domain.,similarity:Contains 1 PNT (pointed) domain.,subunit:Interacts with the DNA-binding domain of the androgen receptor. Interacts with NKX3-1.,tissue specificity:Expressed in a very restricted set of primarily hormone-regulated epithelial tissues with particularly high expression in the prostate gland. Significantly lower expression is seen in other hormone regulated tissues such as mammary gland, salivary gland, and ovary. Expressed in prostate carcinoma cells.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.