

Immunotag™ NLK Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN1253
Product Description	Immunotag™ NLK Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	NLK
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Rat,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from human protein, at AA range: 230-310
Specificity	NLK Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	NLK LAK1
Accession No.	Q9UBE8 O54949 D3ZSZ3

Antibody Specification

Description	<p>catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Activated by tyrosine and threonine phosphorylation (By similarity). Activated by activin.,function:Role in cell fate determination, required for differentiation of bone marrow stromal cells. Acts downstream of MAP3K7 and HIPK2 to negatively regulate the canonical Wnt/beta-catenin signaling pathway and the phosphorylation and destruction of the MYB transcription factor. May suppress a wide range of transcription factors by phosphorylation of the coactivator, CREBBP (By similarity). Involved in TGFbeta-mediated mesoderm induction, acting downstream of MAP3K7/TAK1 to phosphorylate STAT3.,PTM:Dually phosphorylated on Thr-291 and Tyr-293, which activates the enzyme.,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MAP kinase subfamily.,similarity:Contains 1 protein kinase domain.,subcellular location:Predominantly nuclear. A smaller fraction is cytoplasmic.,subunit:Interacts with STAT3 (By similarity). Interacts with RNF138/NARF and TCF7L2/TCF4. Interacts with HIPK2 and MYB.,</p>
Cell Pathway/ Category	MAPK_ERK_Growth,MAPK_G_Protein,WNT,WNT-T CELLAdherens_Junction,
Protein Expression	Amygdala,Placenta,T-cell,Uterus,
Subcellular Localization	intracellular,nucleus,nucleoplasm,cytosol,
Protein Function	<p>catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Activated by tyrosine and threonine phosphorylation (By similarity). Activated by activin.,function:Role in cell fate determination, required for differentiation of bone marrow stromal cells. Acts downstream of MAP3K7 and HIPK2 to negatively regulate the canonical Wnt/beta-catenin signaling pathway and the phosphorylation and destruction of the MYB transcription factor. May suppress a wide range of transcription factors by phosphorylation of the coactivator, CREBBP (By similarity). Involved in TGFbeta-mediated mesoderm induction, acting downstream of MAP3K7/TAK1 to phosphorylate STAT3.,PTM:Dually phosphorylated on Thr-291 and Tyr-293, which activates the enzyme.,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MAP kinase subfamily.,similarity:Contains 1 protein kinase domain.,subcellular location:Predominantly nuclear. A smaller fraction is cytoplasmic.,subunit:Interacts with STAT3 (By similarity). Interacts with RNF138/NARF and TCF7L2/TCF4. Interacts with HIPK2 and MYB.,</p>
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.