

Immunotag™ SL9A2 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN1339
Product Description	Immunotag™ SL9A2 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	SL9A2
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Rat
Host Species	Rabbit
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	SL9A2 Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	SLC9A2 NHE2
Accession No.	Q9UBY0 P48763
Description	solute carrier family 9 member A2(SLC9A2) Homo sapiens This gene encodes a member of the sodium-hydrogen exchanger (NHE) protein family. These proteins are involved in sodium-ion transport by exchanging intracellular hydrogen ions to external sodium ions and help in the regulation of cell pH and volume. The encoded protein is localized to the apical membrane and is involved in apical absorption of sodium. [provided by RefSeq, Jun 2016],

Antibody Specification

Protein Expression	Colon,
Subcellular Localization	plasma membrane,integral component of membrane,
Protein Function	caution:The number, localization and denomination of hydrophobic domains in the Na(+)/H(+) exchangers vary among authors.,function:Involved in pH regulation to eliminate acids generated by active metabolism or to counter adverse environmental conditions. Major proton extruding system driven by the inward sodium ion chemical gradient. Seems to play an important role in colonic sodium absorption.,PTM:Phosphorylated (Possible).,similarity:Belongs to the monovalent cation:proton antiporter 1 (CPA1) transporter (TC 2.A.36) family.,tissue specificity:Expressed in skeletal muscle, colon and kidney. Lower levels in the testis, prostate, ovary, and small intestine.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.