

Immunotag™
 PURA Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN1438
Product Description	Immunotag™ PURA Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PURA
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Rat,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	PURA Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	PURA PUR1
Accession No.	Q00577 P42669 P86252

Antibody Specification

Description	purine rich element binding protein A(PURA) Homo sapiens This gene product is a sequence-specific, single-stranded DNA-binding protein. It binds preferentially to the single strand of the purine-rich element termed PUR, which is present at origins of replication and in gene flanking regions in a variety of eukaryotes from yeasts through humans. Thus, it is implicated in the control of both DNA replication and transcription. Deletion of this gene has been associated with myelodysplastic syndrome and acute myelogenous leukemia. [provided by RefSeq, Jul 2008],
Protein Expression	Brain,Epithelium,Fetal liver,PCR rescued clones,
Subcellular Localization	nuclear chromosome, telomeric region,nucleus,DNA replication factor A complex,cytoplasm,dendrite,neuronal cell body,
Protein Function	function:This is a probable transcription activator that specifically binds the purine-rich single strand of the PUR element located upstream of the MYC gene. May play a role in the initiation of DNA replication and in recombination.,similarity:Belongs to the PUR DNA-binding protein family.,subunit:Homodimer, heterodimer with PURB and heterotrimer with PURB and YBX1/Y-box protein 1.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.