

# Immunotag™ UBE2K Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN1501
Product Description	Immunotag™ UBE2K Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	UBE2K
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	UBE2K Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	UBE2K HIP2 LIG
Accession No.	P61086 P61087

## Antibody Specification

Description	ubiquitin conjugating enzyme E2 K(UBE2K) Homo sapiens The protein encoded by this gene belongs to the ubiquitin-conjugating enzyme family. This protein interacts with RING finger proteins, and it can ubiquitinate huntingtin, the gene product for Huntington's disease. Known functions for this protein include a role in aggregate formation of expanded polyglutamine proteins and the suppression of apoptosis in polyglutamine diseases, a role in the dislocation of newly synthesized MHC class I heavy chains from the endoplasmic reticulum, and involvement in foam cell formation. Multiple transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq, Jul 2008],
Cell Pathway/ Category	Ubiquitin mediated proteolysis,
Protein Expression	Brain,Platelet,Uterus,
Subcellular Localization	nucleus,cytoplasm,filopodium tip,extracellular exosome,
Protein Function	catalytic activity:ATP + ubiquitin + protein lysine = AMP + diphosphate + protein N-ubiquityllysine.,function:Catalyzes the covalent attachment of ubiquitin to other proteins. Mediates the selective degradation of short-lived and abnormal proteins. Ubiquitinates huntingtin. May mediate foam cell formation by the suppression of apoptosis of lipid-bearing macrophages through ubiquitination and subsequent degradation of p53.,induction:By aggregated low-density lipoprotein.,pathway:Protein modification; protein ubiquitination.,PTM:Sumoylation at Lys-14 impairs catalytic activity.,similarity:Belongs to the ubiquitin-conjugating enzyme family.,similarity:Contains 1 UBA domain.,subunit:Interacts with RNF138/NARF.,tissue specificity:Expressed in all tissues tested, including spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocytes, T-lymphocytes, monocytes, granulocytes and bone marrow mononuclear cells. Highly expressed in brain, with highest levels found in cortex and striatum and at lower levels in cerebellum and brainstem.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.