Immunotag[™] VATG3 Polyclonal Antibody

Antibody Specification

Catalog No.	ITN1514
Product Description	Immunotag™ VATG3 Polyclonal Antibody
Size	50 μg, 100 μg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	VATG3
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	VATG3 Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	ATP6V1G3 ATP6G3
Accession No.	Q96LB4 Q8BMC1

Antibody Specification

Description	ATPase H+ transporting V1 subunit G3(ATP6V1G3) Homo sapiens This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'' and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This gene encodes one of three G subunit proteins. Transcript variants encoding
Cell Pathway/ Category	Oxidative phosphorylation, Vibrio cholerae infection, Epithelial cell signaling in Helicobacter pylori infection,
Subcellular Localization	cytosol,plasma membrane,vacuolar proton-transporting V-type ATPase complex,
Protein Function	function:Catalytic subunit of the peripheral V1 complex of vacuolar ATPase (V-ATPase). V- ATPase is responsible for acidifying a variety of intracellular compartments in eukaryotic cells.,similarity:Belongs to the V-ATPase G subunit family.,subunit:V-ATPase is an heteromultimeric enzyme composed of a peripheral catalytic V1 complex (components A to H) attached to an integral membrane V0 proton pore complex (components: a, c, c', c'' and d).,tissue specificity:Kidney.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.

www.gbiosciences.com

© 2018 Geno Technology Inc., USA. All Rights Reserved.