

# Immunotag™ VANG2 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN1517
Product Description	Immunotag™ VANG2 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	VANG2
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Rat,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	VANG2 Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	VANGL2 KIAA1215 STB1
Accession No.	Q9ULK5 Q91ZD4 P84889
Description	VANGL planar cell polarity protein 2(VANGL2) Homo sapiens The protein encoded by this gene is a membrane protein involved in the regulation of planar cell polarity, especially in the stereociliary bundles of the cochlea. The encoded protein transmits directional signals to individual cells or groups of cells in epithelial sheets. This protein is also involved in the development of the neural plate. [provided by RefSeq, Sep 2011],

## Antibody Specification

Cell Pathway/ Category	WNT,WNT-T CELL
Protein Expression	Brain,
Subcellular Localization	stress fiber,plasma membrane,cell-cell junction,integral component of membrane,basolateral plasma membrane,apical plasma membrane,lateral plasma membrane,ER to Golgi transport vesicle,cell pole,
Protein Function	function:Involved in the control of early morphogenesis and patterning of both axial midline structures and the development of neural plate. Plays a role in the regulation of planar cell polarity, particularly in the orientation of stereociliary bundles in the cochlea. Required for polarization and movement of myocardializing cells in the outflow tract and seems to act via RHOA signaling to regulate this process.,similarity:Belongs to the Vang family.,subunit:Interacts through its C-terminal region with the N-terminal half of DVL1, DVL2 and DVL3. The PDZ domain of DVL1, DVL2 and DVL3 is required for the interaction. Also interacts with the PDZ domains of MAGI3, SCRIB/SCRB1 and FZD3.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.