

Immunotag™ TAF1B Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN1630
Product Description	Immunotag™ TAF1B Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	TAF1B
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	TAF1B Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	TAF1B
Accession No.	Q53T94 P97358 D3ZYB7

Antibody Specification

Description	TATA-box binding protein associated factor, RNA polymerase I subunit B(TAF1B) Homo sapiens Initiation of transcription by RNA polymerase I requires the formation of a complex composed of the TATA-binding protein (TBP) and three TBP-associated factors (TAFs) specific for RNA polymerase I. This complex, known as SL1, binds to the core promoter of ribosomal RNA genes to position the polymerase properly and acts as a channel for regulatory signals. This gene encodes one of the SL1-specific TAFs. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2016],
Protein Expression	Colon,Corpus callosum,
Subcellular Localization	nucleus,nucleoplasm,RNA polymerase transcription factor SL1 complex,extracellular exosome,RNA polymerase I core factor complex,
Protein Function	function:Component of the transcription factor SL1/TIFIB complex involved in the assembly of the PIC (preinitiation complex).,similarity:Belongs to the TAF1 family.,subunit:Component of the transcription factor SL1/TIFIB complex which is composed of TBP and at least TAF1A, TAF1B and TAF1C. In the complex, interacts directly with TBP, TAF1A and TAF1C. Interaction of the SL1/TIFIB subunits with TBP excludes interaction with the transcription factor IID (TFIID) subunits. This mutually exclusive binding directs the formation of promoter and RNA polymerase-selective TBP-TAF complexes.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.