

Immunotag™ PLCB1 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN1787
Product Description	Immunotag™ PLCB1 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PLCB1
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	PLCB1 Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	PLCB1 KIAA0581
Accession No.	Q9NQ66 Q9Z1B3 P10687
Description	phospholipase C beta 1(PLCB1) Homo sapiens The protein encoded by this gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of many extracellular signals. This gene is activated by two G-protein alpha subunits, alpha-q and alpha-11. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],

Antibody Specification

Cell Pathway/ Category	Inositol phosphate metabolism,Calcium,Chemokine,Phosphatidylinositol signaling system,Vascular smooth muscle contraction,WNT,WNT-T CELLGap junction,Long-term potentiation,Long-term depression,GnRH,Melanogenesis,Alzheimer's disease,Huntington's disease,
Protein Expression	Brain,Epithelium,Placenta,Testis,
Subcellular Localization	nuclear chromatin,nucleus,cytoplasm,cytosol,nuclear speck,nuclear membrane,myelin sheath,extracellular exosome,
Protein Function	<p>catalytic activity:1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate + H(2)O = 1D-myo-inositol 1,4,5-trisphosphate + diacylglycerol.,cofactor:Calcium.,function:The production of the second messenger molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes.,miscellaneous:The receptor-mediated activation of PLC-beta-1 is mediated by two G-protein alpha subunits, alpha-Q and alpha-11.,similarity:Contains 1 C2 domain.,similarity:Contains 1 PI-PLC X-box domain.,similarity:Contains 1 PI-PLC Y-box domain.,</p>
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.