

# Immunotag™ PCP Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN1985
Product Description	Immunotag™ PCP Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PCP
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	PCP Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	PRCP PCP
Accession No.	P42785 Q7TMR0

## Antibody Specification

Description	prolylcarboxypeptidase(PRCP) Homo sapiens This gene encodes a member of the peptidase S28 family of serine exopeptidases. The encoded preproprotein is proteolytically processed to generate the mature lysosomal prolylcarboxypeptidase. This enzyme cleaves C-terminal amino acids linked to proline in peptides such as angiotension II, III and des-Arg9-bradykinin. The cleavage occurs at acidic pH, but the enzyme activity is retained with some substrates at neutral pH. This enzyme has been shown to be an activator of the cell matrix-associated prekallikrein. The importance of angiotension II, one of the substrates of this enzyme, in regulating blood pressure and electrolyte balance suggests that this gene may be related to essential hypertension. A pseudogene of this gene has been identified on chromosome 2. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [pr
Protein Expression	Brain,Kidney,Liver,Lung,
Subcellular Localization	lysosome,plasma membrane,basal part of cell,extracellular exosome,
Protein Function	catalytic activity:Cleavage of a -Pro- -Xaa bond to release a C-terminal amino acid.,function:Cleaves C-terminal amino acids linked to proline in peptides such as angiotensin II, III and des-Arg9-bradykinin. This cleavage occurs at acidic pH, but enzymatic activity is retained with some substrates at neutral pH.,similarity:Belongs to the peptidase S28 family.,subunit:Homodimer.,tissue specificity:Highest levels in placenta, lung and liver. Also present in heart, brain, pancreas and kidney.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.