

Immunotag™ TXND2 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN2078
Product Description	Immunotag™ TXND2 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	TXND2
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Rat,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	TXND2 Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	TXNDC2 SPTRX SPTRX1
Accession No.	Q86VQ3 Q6P902 Q5XHX6

Antibody Specification

Description	function:Probably plays a regulatory role in sperm development. May participate in regulation of fibrous sheath (FS) assembly by supporting the formation of disulfide bonds during sperm tail morphogenesis. May also be required to rectify incorrect disulfide pairing and generate suitable pairs between the FS constituents. Can reduce disulfide bonds in vitro in the presence of NADP and thioredoxin reductase.,similarity:Contains 1 thioredoxin domain.,subcellular location:In ejaculated spermatozoa, it localizes in the caudal region of the head to the end of the principal piece.,tissue specificity:Testis-specific. Only expressed during spermiogenesis, prominently in round and elongating spermatids.,
Protein Expression	Brain,Testis,
Subcellular Localization	outer dense fiber,nucleus,nucleolus,cytoplasm,mitochondrion,
Protein Function	function:Probably plays a regulatory role in sperm development. May participate in regulation of fibrous sheath (FS) assembly by supporting the formation of disulfide bonds during sperm tail morphogenesis. May also be required to rectify incorrect disulfide pairing and generate suitable pairs between the FS constituents. Can reduce disulfide bonds in vitro in the presence of NADP and thioredoxin reductase.,similarity:Contains 1 thioredoxin domain.,subcellular location:In ejaculated spermatozoa, it localizes in the caudal region of the head to the end of the principal piece.,tissue specificity:Testis-specific. Only expressed during spermiogenesis, prominently in round and elongating spermatids.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.