

Immunotag™ PPM1B Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN2285
Product Description	Immunotag™ PPM1B Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PPM1B
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Rat,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from human protein, at AA range: 310-390
Specificity	PPM1B Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	PPM1B PP2CB
Accession No.	O75688 P36993 P35815

Antibody Specification

Description	protein phosphatase, Mg ²⁺ /Mn ²⁺ dependent 1B (PPM1B) Homo sapiens The protein encoded by this gene is a member of the PP2C family of Ser/Thr protein phosphatases. PP2C family members are known to be negative regulators of cell stress response pathways. This phosphatase has been shown to dephosphorylate cyclin-dependent kinases (CDKs), and thus may be involved in cell cycle control. Overexpression of this phosphatase is reported to cause cell-growth arrest or cell death. Alternative splicing results in multiple transcript variants encoding different isoforms. Additional transcript variants have been described, but currently do not represent full-length sequences. [provided by RefSeq, Jul 2008],
Cell Pathway/ Category	MAPK_ERK_Growth,MAPK_G_Protein,
Protein Expression	Adrenal gland,Bladder,Brain,Liver,Stomach,
Subcellular Localization	cytosol,membrane,
Protein Function	Additional isoforms seem to exist,catalytic activity:A phosphoprotein + H ₂ O = a protein + phosphate.,cofactor:Binds 2 magnesium or manganese ions per subunit.,function:Enzyme with a broad specificity. Dephosphorylates CDK2 and CDK6 in vitro.,similarity:Belongs to the PP2C family.,subunit:Monomer.,tissue specificity:Highly expressed in heart and skeletal muscle.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.