

Immunotag™ SMAGP Polyclonal Antibody

| Antibody Specification | |
|------------------------|--|
| Catalog No. | ITN2314 |
| Product Description | Immunotag™ SMAGP Polyclonal Antibody |
| Size | 50 µg, 100 µg |
| Conjugation | HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647 |
| IMPORTANT NOTE | This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return. |
| Target Protein | SMAGP |
| Clonality | Polyclonal |
| Storage/Stability | -20°C/1 year |
| Application | WB,ELISA |
| Recommended Dilution | WB 1:500-2000 ELISA 1:5000-20000 |
| Concentration | 1 mg/ml |
| Reactive Species | Human,Rat,Mouse |
| Host Species | Rabbit |
| Immunogen | Synthesized peptide derived from human protein, at AA range: 30-110 |
| Specificity | SMAGP Polyclonal Antibody detects endogenous levels of protein. |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen |
| Form | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Gene Name | SMAGP |
| Accession No. | Q0VAQ4 Q99KC7 Q7TPF1 |

Antibody Specification

| | |
|--------------------------|--|
| Description | function:May play a role in epithelial cell-cell contacts. May play a role in tumor invasiveness and metastasis formation.,PTM:O-glycosylated. The O-glycan is modified with sialic acid residues.,similarity:Belongs to the SMAGP family.,subcellular location:Predominantly on lateral parts of the membrane, at cell-cell epithelial junctions. Detected on cytoplasmic membranes in undifferentiated tumors.,tissue specificity:Detected in breast, endometrium, colon and biliary tract. Detected in polarized epithelial structures characterized by cell-cell adhesion (at protein level)., |
| Protein Expression | Colon, |
| Subcellular Localization | nucleoplasm,plasma membrane,integral component of plasma membrane,cell-cell adherens junction,integral component of membrane,cell junction,cytoplasmic vesicle membrane, |
| Protein Function | function:May play a role in epithelial cell-cell contacts. May play a role in tumor invasiveness and metastasis formation.,PTM:O-glycosylated. The O-glycan is modified with sialic acid residues.,similarity:Belongs to the SMAGP family.,subcellular location:Predominantly on lateral parts of the membrane, at cell-cell epithelial junctions. Detected on cytoplasmic membranes in undifferentiated tumors.,tissue specificity:Detected in breast, endometrium, colon and biliary tract. Detected in polarized epithelial structures characterized by cell-cell adhesion (at protein level)., |
| Usage | For Research Use Only! Not for diagnostic or therapeutic procedures. |