

Immunotag™ SPON1 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN2408
Product Description	Immunotag™ SPON1 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	SPON1
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Rat,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from human protein, at AA range: 210-290
Specificity	SPON1 Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	SPON1 KIAA0762 VSGP
Accession No.	Q9HCB6 Q8VCC9 P35446

Antibody Specification

Description	function:Cell adhesion protein that promotes the attachment of spinal cord and sensory neuron cells and the outgrowth of neurites in vitro. May contribute to the growth and guidance of axons in both the spinal cord and the PNS (By similarity). Major factor for vascular smooth muscle cell.,similarity:Contains 1 reelin domain.,similarity:Contains 1 spondin domain.,similarity:Contains 6 TSP type-1 domains.,subunit:Binds to the central extracellular domain of APP and inhibits beta-secretase cleavage of APP.,tissue specificity:Highest expression in lung, lower expression in brain, heart, kidney, liver and testis, and lowest expression in pancreas, skeletal muscle and ovary. Not expressed in spleen.,
Protein Expression	Brain,Ovary,Placenta,Pooled,
Subcellular Localization	proteinaceous extracellular matrix,extracellular space,endoplasmic reticulum lumen,extracellular matrix,
Protein Function	function:Cell adhesion protein that promotes the attachment of spinal cord and sensory neuron cells and the outgrowth of neurites in vitro. May contribute to the growth and guidance of axons in both the spinal cord and the PNS (By similarity). Major factor for vascular smooth muscle cell.,similarity:Contains 1 reelin domain.,similarity:Contains 1 spondin domain.,similarity:Contains 6 TSP type-1 domains.,subunit:Binds to the central extracellular domain of APP and inhibits beta-secretase cleavage of APP.,tissue specificity:Highest expression in lung, lower expression in brain, heart, kidney, liver and testis, and lowest expression in pancreas, skeletal muscle and ovary. Not expressed in spleen.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.