

Immunotag™ HIC2 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN2963
Product Description	Immunotag™ HIC2 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	HIC2
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	HIC2 Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	HIC2 HRG22 KIAA1020 ZBTB30
Accession No.	Q96JB3 Q9JLZ6
Description	function:Transcriptional repressor.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family. Hic subfamily.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 5 C2H2-type zinc fingers.,subunit:Interacts with CtBP.,tissue specificity:Highest levels in cerebellum.,
Protein Expression	Brain,Placenta,Testis,

Antibody Specification

Subcellular Localization	nucleus,
Protein Function	function:Transcriptional repressor.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family. Hic subfamily.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 5 C2H2-type zinc fingers.,subunit:Interacts with CtBP.,tissue specificity:Highest levels in cerebellum.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.