Immunotag™ Rsk-1/2/3/4 (phospho Ser221/227/S218/232) Polyclonal Antibody

| Antibody Specification | |
|-------------------------|--|
| Catalog No. | ITP0436 |
| Product Description | Immunotag™ Rsk-1/2/3/4 (phospho Ser221/227/S218/232) Polyclonal Antibody |
| Size | 50 μg, 100 μg |
| Conjugation | HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647 |
| IMPORTANT NOTE | This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return. |
| Target Protein | Rsk-1/2/3/4 |
| Clonality | Polyclonal |
| Storage/Stability | -20°C/1 year |
| Application | WB,ELISA |
| Recommended Dilution | Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. |
| Concentration | 1 mg/ml |
| Reactive Species | Human, Mouse |
| Host Species | Rabbit |
| Immunogen | The antiserum was produced against synthesized peptide derived from human RSK1/2/3/4 around the phosphorylation site of Ser221/227/S218/232. AA range:191-240 |
| Specificity | Phospho-Rsk-1/2/3/4 (S221/227/S218/232) Polyclonal Antibody detects endogenous levels of Rsk-1/2/3/4 protein only when phosphorylated at S221/227/S218/232. |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen |
| Form | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Gene Name | RPS6KA1 |
| Accession No. | Q15418/P51812/Q15349/Q9UK32 |
| Alternate Names | RPS6KA1; MAPKAPK1A; RSK1; Ribosomal protein S6 kinase alpha-1; S6K-alpha-1; 90 kDa ribosomal protein S6 kinase 1; p90-RSK 1; p90RSK1; p90S6K; MAP kinase-activated protein kinase 1a; MAPK-activated protein kinase 1a; MAPKAP |

| Antibody Specification | |
|-----------------------------|--|
| Description | ribosomal protein S6 kinase A1(RPS6KA1) Homo sapiens This gene encodes a member of the RSK (ribosomal S6 kinase) family of serine/threonine kinases. This kinase contains 2 nonidentical kinase catalytic domains and phosphorylates various substrates, including members of the mitogen-activated kinase (MAPK) signalling pathway. The activity of this protein has been implicated in controlling cell growth and differentiation. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008], |
| Cell Pathway/ Category | Regulates Angiogenesis, Insulin Receptor, B Cell Receptor, AMPK |
| Protein Expression | Colon,Epithelium, |
| Subcellular Localization | nucleus,nucleoplasm,cytoplasm,cytosol, |
| Protein Function | catalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,cofactor:Magnesium.,enzyme regulation:Activated by multiple phosphorylations on threonine and serine residues.,function:Serine/threonine kinase that may play a role in mediating the growth-factor and stress induced activation of the transcription factor CREB.,PTM:Autophosphorylated on Ser-380, as part of the activation process.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase family. S6 kinase subfamily.,similarity:Contains 1 AGC-kinase C-terminal domain.,similarity:Contains 2 protein kinase domains.,subunit:Forms a complex with either ERK1 or ERK2 in quiescent cells. Transiently dissociates following mitogenic stimulation., |
| Usage | For Research Use Only! Not for diagnostic or therapeutic procedures. |

www.gbiosciences.com

© 2018 Geno Technology Inc., USA. All Rights Reserved.