

Immunotag™ PLC γ2 (phospho Tyr753) Polyclonal Antibody

Antibody Specification	
Catalog No.	ITP0608
Product Description	Immunotag™ PLC γ2 (phospho Tyr753) Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PLC γ2 (Tyr753)
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	Synthesized phospho-peptide around the phosphorylation site of human PLC γ2 (phospho Tyr753)
Specificity	Phospho-PLC γ2 (Y753) Polyclonal Antibody detects endogenous levels of PLC γ2 protein only when phosphorylated at Y753.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	PLCG2
Accession No.	P16885 Q8CIH5 P24135
Alternate Names	PLCG2; 1-phosphatidylinositol 4; 5-bisphosphate phosphodiesterase gamma-2; Phosphoinositide phospholipase C-gamma-2; Phospholipase C-IV; PLC-IV; Phospholipase C-gamma-2; PLC-gamma-2

Antibody Specification

Description	phospholipase C gamma 2(PLCG2) Homo sapiens The protein encoded by this gene is a transmembrane signaling enzyme that catalyzes the conversion of 1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate to 1D-myo-inositol 1,4,5-trisphosphate (IP3) and diacylglycerol (DAG) using calcium as a cofactor. IP3 and DAG are second messenger molecules important for transmitting signals from growth factor receptors and immune system receptors across the cell membrane. Mutations in this gene have been found in autoinflammation, antibody deficiency, and immune dysregulation syndrome and familial cold autoinflammatory syndrome 3. [provided by RefSeq, Mar 2014],
Cell Pathway/ Category	Inositol phosphate metabolism,ErbB_HER,Calcium,Phosphatidylinositol signaling system,VEGF,Natural killer cell mediated cytotoxicity,B_Cell_Antigen,Fc epsilon RI,Fc gamma R-mediated phagocytosis,Leukocyte transendothelial migration,Neurotrophin,Vibrio cholerae infection,Epithelial cell signaling in Helicobacter pylori infection,Pathways in cancer,Glioma,Non-small cell lung cancer,
Protein Expression	Lymph,Lymphoblast,Spleen,T-cell,
Subcellular Localization	intracellular,cytosol,plasma membrane,extracellular exosome,
Protein Function	catalytic activity:1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate + H(2)O = 1D-myo-inositol 1,4,5-trisphosphate + diacylglycerol.,cofactor:Calcium.,function:The production of the second messenger molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes. It is a crucial enzyme in transmembrane signaling.,PTM:Phosphorylated on tyrosine residues; upon ligand-induced activation of a variety of growth factor receptors and immune system receptors. Increases phospholipase activity.,similarity:Contains 1 C2 domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 PI-PLC X-box domain.,similarity:Contains 1 PI-PLC Y-box domain.,similarity:Contains 1 SH3 domain.,similarity:Contains 2 SH2 domains.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.