

Immunotag™ CD30-L Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT0750
Product Description	Immunotag™ CD30-L Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	CD30-L
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human CD153. AA range:71-120
Specificity	CD30-L Polyclonal Antibody detects endogenous levels of CD30-L protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	TNFSF8
Accession No.	P32971 P32972
Alternate Names	TNFSF8; CD30L; CD30LG; Tumor necrosis factor ligand superfamily member 8; CD30 ligand; CD30-L; CD antigen CD153

Antibody Specification

Description	tumor necrosis factor superfamily member 8(TNFSF8) Homo sapiens The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF8/CD30, which is a cell surface antigen and a marker for Hodgkin lymphoma and related hematologic malignancies. The engagement of this cytokine expressed on B cell surface plays an inhibitory role in modulating Ig class switch. This cytokine was shown to enhance cell proliferation of some lymphoma cell lines, while to induce cell death and reduce cell proliferation of other lymphoma cell lines. The pleiotropic biologic activities of this cytokine on different CD30+ lymphoma cell lines may play a pathophysiologic role in Hodgkin's and some non-Hodgkin's lymphomas. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2011],
Cell Pathway/ Category	Cytokine-cytokine receptor interaction,
Protein Expression	Brain,Thymus,
Subcellular Localization	extracellular space,integral component of plasma membrane,integral component of membrane,
Protein Function	function:Cytokine that binds to TNFRSF8/CD30. Induces proliferation of T-cells.,similarity:Belongs to the tumor necrosis factor family.,subunit:Homotrimer.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.