

Immunotag™ GPR116 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT1961
Product Description	Immunotag™ GPR116 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	GPR116
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	IHC-p,IF,ELISA
Recommended Dilution	Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human GPR116. AA range:1121-1170
Specificity	GPR116 Polyclonal Antibody detects endogenous levels of GPR116 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	GPR116
Accession No.	Q8IZF2
Alternate Names	GPR116; KIAA0758; Probable G-protein coupled receptor 116

Antibody Specification

Description	function:May have a role in the regulation of acid-base balance.,PTM:Proteolytically cleaved into 2 highly conserved sites: one in the SEA domain and the other in the stalk domain region preceding the first transmembrane. The later 2 subunits, the extracellular subunit and the seven-transmembrane subunit, remain tightly associated and non-covalently linked.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Belongs to the G-protein coupled receptor 2 family. LN-TM7 subfamily.,similarity:Contains 1 GPS domain.,similarity:Contains 1 SEA domain.,similarity:Contains 3 Ig-like (immunoglobulin-like) domains.,subunit:Exists as disulfide-linked dimers at the cell surface.,
Protein Expression	Brain,Plasma,Testis,
Subcellular Localization	plasma membrane,cell surface,integral component of membrane,cytoplasmic vesicle,apical part of cell,
Protein Function	function:May have a role in the regulation of acid-base balance.,PTM:Proteolytically cleaved into 2 highly conserved sites: one in the SEA domain and the other in the stalk domain region preceding the first transmembrane. The later 2 subunits, the extracellular subunit and the seven-transmembrane subunit, remain tightly associated and non-covalently linked.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Belongs to the G-protein coupled receptor 2 family. LN-TM7 subfamily.,similarity:Contains 1 GPS domain.,similarity:Contains 1 SEA domain.,similarity:Contains 3 Ig-like (immunoglobulin-like) domains.,subunit:Exists as disulfide-linked dimers at the cell surface.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.