

Immunotag™ GPR120 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT1964
Product Description	Immunotag™ GPR120 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	GPR120
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	Synthesized peptide derived from GPR120, at AA range: 190-270
Specificity	GPR120 Polyclonal Antibody detects endogenous levels of GPR120 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	O3FAR1
Accession No.	Q5NUL3 Q7TMA4 Q2AC31
Alternate Names	O3FAR1; GPR120; GPR129; PGR4; Omega-3 fatty acid receptor 1; G-protein coupled receptor 120; G-protein coupled receptor 129; G-protein coupled receptor GT01; G-protein coupled receptor PGR4

Antibody Specification

Description	free fatty acid receptor 4(FFAR4) Homo sapiens This gene encodes a G protein-coupled receptor (GPR) which belongs to the rhodopsin family of GPRs. The encoded protein functions as a receptor for free fatty acids, including omega-3, and participates in suppressing anti-inflammatory responses and insulin sensitizing. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Feb 2012],
Subcellular Localization	plasma membrane,integral component of plasma membrane,integral component of membrane,endocytic vesicle,
Protein Function	function:Receptor for unsaturated long-chain free fatty acid (FAA). Binding of the ligand promotes the secretion of glucagon-like peptide-1 from the gastro-intestinal tract.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Abundant expression in the intestinal tract.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.