

Immunotag™ GPR18 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT2005
Product Description	Immunotag™ GPR18 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	GPR18
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,IF,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	Synthesized peptide derived from GPR18, at AA range: 160-240
Specificity	GPR18 Polyclonal Antibody detects endogenous levels of GPR18 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	GPR18
Accession No.	Q14330 Q8K1Z6 A1A5S3
Alternate Names	GPR18; GPCRW; N-arachidonyl glycine receptor; NAGly receptor; G-protein coupled receptor 18

Antibody Specification

Description	function:Receptor for N-arachidonyl glycine. The activity of this receptor is mediated by G proteins which inhibit adenylyl cyclase. May contribute to regulation of the immune system.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Most abundant in testis and spleen. Highly expressed in CD4 and CD8-positive T-cells as well as CD19-positive B-cells.,
Protein Expression	Dendritic cell,Lymph,Testis,
Subcellular Localization	plasma membrane,integral component of plasma membrane,integral component of membrane,
Protein Function	function:Receptor for N-arachidonyl glycine. The activity of this receptor is mediated by G proteins which inhibit adenylyl cyclase. May contribute to regulation of the immune system.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Most abundant in testis and spleen. Highly expressed in CD4 and CD8-positive T-cells as well as CD19-positive B-cells.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.