

Immunotag™ GPR41 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT2020
Product Description	Immunotag™ GPR41 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	GPR41
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human FFAR3. AA range:11-60
Specificity	GPR41 Polyclonal Antibody detects endogenous levels of GPR41 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	FFAR3
Accession No.	O14843 Q3UFD7
Alternate Names	FFAR3; GPR41; Free fatty acid receptor 3; G-protein coupled receptor 41
Description	function:Receptor for short chain fatty acids through a G(i)-protein-mediated inhibition of adenylyl cyclase and elevation of intracellular calcium. The rank order of potency for agonists of this receptor is propionate = pentanoate = butyrate > acetate > formate.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Highest level in adipose tissue, and lower expression across all tissues tested.,

Antibody Specification

Protein Expression	Brain,Ovary,
Subcellular Localization	plasma membrane,integral component of plasma membrane,integral component of membrane,
Protein Function	function:Receptor for short chain fatty acids through a G(i)-protein-mediated inhibition of adenylyl cyclase and elevation of intracellular calcium. The rank order of potency for agonists of this receptor is propionate = pentanoate = butyrate > acetate > formate.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Highest level in adipose tissue, and lower expression across all tissues tested.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.