

Immunotag™ Hint1 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT2136
Product Description	Immunotag™ Hint1 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Hint1
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,IF,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	Synthesized peptide derived from Hint1, at AA range: 40-120
Specificity	Hint1 Polyclonal Antibody detects endogenous levels of Hint1 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	HINT1
Accession No.	P49773 P70349 P62959
Alternate Names	HINT1; HINT; PKCI1; PRKCNH1; Histidine triad nucleotide-binding protein 1; Adenosine 5'-monophosphoramidase; Protein kinase C inhibitor 1; Protein kinase C-interacting protein 1; PKCI-1

Antibody Specification

Description	histidine triad nucleotide binding protein 1(HINT1) Homo sapiens This gene encodes a protein that hydrolyzes purine nucleotide phosphoramidates substrates, including AMP-morpholidate, AMP-N-alanine methyl ester, AMP-alpha-acetyl lysine methyl ester, and AMP-NH2. The encoded protein interacts with these substrates via a histidine triad motif. This gene is considered a tumor suppressor gene. In addition, mutations in this gene can cause autosomal recessive neuromyotonia and axonal neuropathy. There are several related pseudogenes on chromosome 7. Several transcript variants have been observed. [provided by RefSeq, Dec 2015],
Protein Expression	Cervix,Platelet,Prostate,
Subcellular Localization	histone deacetylase complex,nucleus,cytoplasm,cytoskeleton,plasma membrane,extracellular exosome,
Protein Function	caution:Was originally thought to be a protein kinase C inhibitor and to bind zinc in solution. Both seem to be incorrect.,domain:The histidine triad, also called HIT motif, forms part of the binding loop for the alpha-phosphate of purine mononucleotide.,function:Hydrolyzes adenosine 5'-monophosphoramidate substrates such as AMP-morpholidate, AMP-N-alanine methyl ester, AMP-alpha-acetyl lysine methyl ester and AMP-NH2.,similarity:Belongs to the HINT family.,similarity:Contains 1 HIT domain.,subcellular location:Interaction with CDK7 leads to a more nuclear localization.,subunit:Homodimer. Interacts with CDK7.,tissue specificity:Widely expressed.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.