

# Immunotag™ MRP-S22 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT2881
Product Description	Immunotag™ MRP-S22 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	MRPS22
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,IF,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Monkey
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human MRPS22. AA range:231-280
Specificity	MRP-S22 Polyclonal Antibody detects endogenous levels of MRP-S22 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	MRPS22
Accession No.	P82650 Q9CXW2
Alternate Names	MRPS22; C3orf5; RPMS22; GK002; 28S ribosomal protein S22; mitochondrial; MRP-S22; S22mt

## Antibody Specification

Description	mitochondrial ribosomal protein S22(MRPS22) Homo sapiens Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein that does not seem to have a counterpart in prokaryotic and fungal-mitochondrial ribosomes. This gene lies telomeric of and is transcribed in the opposite direction from the forkhead box L2 gene. A pseudogene
Protein Expression	Brain,Liver cancer,Muscle,Placenta,
Subcellular Localization	mitochondrion,mitochondrial inner membrane,mitochondrial ribosome,mitochondrial small ribosomal subunit,ribosome,
Protein Function	disease:Defects in MRPS22 are the cause of combined oxidative phosphorylation deficiency type 5 (COXPD5) [MIM:611719]. COXPD5 is an antenatal mitochondrial disease. Patients show edema, cardiomyopathy, tubulopathy, and hypotonia.,subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.