Immunotag[™] MRP-S27 Polyclonal Antibody

| Antibody Specification | | |
|------------------------|--|--|
| Catalog No. | ITT2884 | |
| Product Description | Immunotag™ MRP-S27 Polyclonal Antibody | |
| Size | 50 μg, 100 μg | |
| Conjugation | HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647 | |
| IMPORTANT NOTE | This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return. | |
| Target Protein | MRPS27 | |
| Clonality | Polyclonal | |
| Storage/Stability | -20°C/1 year | |
| Application | WB,ELISA | |
| Recommended Dilution | Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. | |
| Concentration | 1 mg/ml | |
| Reactive Species | Human | |
| Host Species | Rabbit | |
| Immunogen | The antiserum was produced against synthesized peptide derived from human MRPS27. AA range:16-65 | |
| Specificity | MRP-S27 Polyclonal Antibody detects endogenous levels of MRP-S27 protein. | |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen | |
| Form | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. | |
| Gene Name | MRPS27 | |
| Accession No. | Q92552 Q8BK72 | |
| Alternate Names | MRPS27; KIAA0264; 28S ribosomal protein S27; mitochondrial; MRP-S27; S27mt | |

| Antibody Specification | |
|-----------------------------|---|
| Description | mitochondrial ribosomal protein S27(MRPS27) Homo sapiens Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein that may be a functional partner of the death associated protein 3 (DAP3). Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Nov 2013], |
| Protein Expression | Bone marrow,Colon,Lung,Skin,Uterus, |
| Subcellular Localization | mitochondrion,mitochondrial inner membrane,ribosome, |
| Protein Function | similarity:Contains 2 PPR (pentatricopeptide) repeats.,subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., |
| Usage | For Research Use Only! Not for diagnostic or therapeutic procedures. |

www.gbiosciences.com

© 2018 Geno Technology Inc., USA. All Rights Reserved.