

# Immunotag™ MYLIP Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT2939
Product Description	Immunotag™ MYLIP Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	MYLIP
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	Synthesized peptide derived from MYLIP, at AA range: 130-210
Specificity	MYLIP Polyclonal Antibody detects endogenous levels of MYLIP protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	MYLIP
Accession No.	Q8WY64 Q8BM54 D3ZDI6
Alternate Names	MYLIP; BZF1; IDOL; BM-023; PP5242; E3 ubiquitin-protein ligase MYLIP; Inducible degrader of the LDL-receptor; Idol; Myosin regulatory light chain interacting protein; MIR

## Antibody Specification

Description	myosin regulatory light chain interacting protein(MYLIP) Homo sapiens The ERM protein family members ezrin, radixin, and moesin are cytoskeletal effector proteins linking actin to membrane-bound proteins at the cell surface. Myosin regulatory light chain interacting protein (MYLIP) is a novel ERM-like protein that interacts with myosin regulatory light chain and inhibits neurite outgrowth. [provided by RefSeq, Jul 2008],
Protein Expression	Bone marrow,Brain,Placenta,
Subcellular Localization	intracellular,cytosol,cytoskeleton,plasma membrane,extrinsic component of membrane,
Protein Function	developmental stage:Expressed in fetal tissues and higher levels were detected in placenta and fetal lung.,domain:The RING domain mediates ubiquitination and the neurite outgrowth inhibitory activity.,function:E3 ubiquitin-protein ligase that mediates ubiquitination and subsequent proteasomal degradation of myosin regulatory light chain (MRLC). Stable or transient overexpression of MIR induces proteasomal degradation of MRLC inhibiting neurite outgrowth in presence of NGF. MIR counteracts the stabilization of MRLC by MIR-interacting saposin-like protein (MSAP/TMEM4) and reduces MSAP-stimulated neurite outgrowth.,pathway:Protein modification; protein ubiquitination.,PTM:Autoubiquitinated.,similarity:Contains 1 FERM domain.,similarity:Contains 1 RING-type zinc finger.,subunit:Interacts with myosin regulatory light chain (MRLC) and TMEM4.,tissue specificity:Ubiquitously expressed.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.