Immunotag[™] Na+/K+-ATPase α1 Polyclonal Antibody

| Antibody Specification | |
|-------------------------|--|
| Catalog No. | ITT2972 |
| Product Description | Immunotag™ Na+/K+-ATPase α1 Polyclonal Antibody |
| Size | 50 μg, 100 μg |
| Conjugation | HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647 |
| IMPORTANT NOTE | This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return. |
| Target Protein | Na+/K+-ATPase α1 |
| Clonality | Polyclonal |
| Storage/Stability | -20°C/1 year |
| Application | WB,IF,ELISA |
| Recommended Dilution | Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. |
| Concentration | 1 mg/ml |
| Reactive Species | Rat |
| Host Species | Rabbit |
| Immunogen | Synthesized peptide derived from Na+/K+-ATPase α 1, at AA range: 10-90 |
| Specificity | Na+/K+-ATPase $\alpha 1$ Polyclonal Antibody detects endogenous levels of Na+/K+-ATPase $\alpha 1$ protein. |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen |
| Form | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Gene Name | ATP1A1 |
| Accession No. | P06685 |
| Alternate Names | ATP1A1; Sodium/potassium-transporting ATPase subunit alpha-1; Na(+)/K(+) ATPase alpha-1 subunit; Sodium pump subunit alpha-1 |

| Antibody Specification | | |
|------------------------|---|--|
| Description | The ATPase Na+/K+ transporting subunit alpha 1 encoded by ATP1A1 belongs to the family of P-type cation transport ATPases, and to the subfamily of Na+/K+ -ATPases. Na+/K+ -ATPase is an integral membrane protein responsible for establishing and maintaining the electrochemical gradients of Na and K ions across the plasma membrane. These gradients are essential for osmoregulation, for sodium-coupled transport of a variety of organic and inorganic molecules, and for electrical excitability of nerve and muscle. This enzyme is composed of two subunits, a large catalytic subunit (alpha) and a smaller glycoprotein subunit (beta). The catalytic subunit of Na+/K+ -ATPase is encoded by multiple genes. This gene encodes an alpha 1 subunit. Multiple transcript variants encoding different isoforms have been found for this gene. | |
| Usage | For Research Use Only! Not for diagnostic or therapeutic procedures. | |

www.gbiosciences.com

© 2018 Geno Technology Inc., USA. All Rights Reserved.