

Immunotag™ Nectin 3 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT3024
Product Description	Immunotag™ Nectin 3 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Nectin 3
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human PVRL3. AA range:184-233
Specificity	Nectin 3 Polyclonal Antibody detects endogenous levels of Nectin 3 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	PVRL3
Accession No.	Q9NQS3 Q9JLB9
Alternate Names	PVRL3; PRR3; Poliovirus receptor-related protein 3; CDw113; Nectin-3; CD antigen CD113

Antibody Specification

Description	nectin cell adhesion molecule 3(NECTIN3) Homo sapiens This gene encodes a member of the nectin family of proteins, which function as adhesion molecules at adherens junctions. This family member interacts with other nectin-like proteins and with afadin, a filamentous actin-binding protein involved in the regulation of directional motility, cell proliferation and survival. This gene plays a role in ocular development involving the ciliary body. Mutations in this gene are believed to result in congenital ocular defects. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011],
Cell Pathway/ Category	Cell adhesion molecules (CAMs),Adherens_Junction,
Protein Expression	Brain,Cervix,Fetal kidney,Kidney,Placenta,
Subcellular Localization	plasma membrane,integral component of plasma membrane,cell-cell junction,cell-cell adherens junction,apical junction complex,cell-cell contact zone,postsynaptic membrane,
Protein Function	function:Plays a role in cell-cell adhesion through heterophilic trans-interactions with nectin-like proteins or nectins, such as trans-interaction with PVRL2/nectin-2 at Sertoli-spermatid junctions. Trans-interaction with PVR induces activation of CDC42 and RAC small G proteins through common signaling molecules such as SRC and RAP1. Also involved in the formation of cell-cell junctions, including adherens junctions and synapses. Induces endocytosis-mediated down-regulation of PVR from the cell surface, resulting in reduction of cell movement and proliferation. Plays a role in the morphology of the ciliary body.,similarity:Belongs to the nectin family.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,subunit:Cis- and trans-homodimer. Can form trans-heterodimers with PVRL1/nectin-1, PVRL2/nectin-2, PVR, IGSF4B/Necl-1 and with IGSF4. Interacts with MLLT4/afadin. Binds with low affinity to TIGIT.,tissue specificity:Predominantly expressed in testis and placenta as well as in many cell lines, including epithelial cell lines.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.