

Immunotag™ NF-YB Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT3091
Product Description	Immunotag™ NF-YB Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	NFYB
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	Synthesized peptide derived from NF-YB, at AA range: 10-90
Specificity	NF-YB Polyclonal Antibody detects endogenous levels of NF-YB protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	NFYB
Accession No.	P25208 P63139 P63140
Alternate Names	NFYB; HAP3; Nuclear transcription factor Y subunit beta; CAAT box DNA-binding protein subunit B; Nuclear transcription factor Y subunit B; NF-YB

Antibody Specification

Description	nuclear transcription factor Y subunit beta(NFYB) Homo sapiens The protein encoded by this gene is one subunit of a trimeric complex, forming a highly conserved transcription factor that binds with high specificity to CCAAT motifs in the promoter regions in a variety of genes. This gene product, subunit B, forms a tight dimer with the C subunit, a prerequisite for subunit A association. The resulting trimer binds to DNA with high specificity and affinity. Subunits B and C each contain a histone-like motif. Observation of the histone nature of these subunits is supported by two types of evidence; protein sequence alignments and experiments with mutants. [provided by RefSeq, Jul 2008],
Cell Pathway/ Category	Antigen processing and presentation,
Protein Expression	Urinary bladder,
Subcellular Localization	nucleus,nucleoplasm,CCAAT-binding factor complex,protein-DNA complex,
Protein Function	domain:Can be divided into 3 domains: the weakly conserved A domain, the highly conserved B domain thought to be involved in subunit interaction and DNA binding, and the Glu-rich C domain.,function:Stimulates the transcription of various genes by recognizing and binding to a CCAAT motif in promoters, for example in type 1 collagen, albumin and beta-actin genes.,similarity:Belongs to the NFYB/HAP3 subunit family.,subunit:Heterotrimeric transcription factor composed of three components, NF-YA, NF-YB and NF-YC. NF-YB and NF-YC must interact and dimerize for NF-YA association and DNA binding.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.