

Immunotag™ NHE-7 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT3118
Product Description	Immunotag™ NHE-7 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	NHE-7
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human SLC9A7. AA range:531-580
Specificity	NHE-7 Polyclonal Antibody detects endogenous levels of NHE-7 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	SLC9A7
Accession No.	Q96T83 Q8BLV3
Alternate Names	SLC9A7; NHE7; Sodium/hydrogen exchanger 7; Na(+)/H(+) exchanger 7; NHE-7; Solute carrier family 9 member 7

Antibody Specification

Description	solute carrier family 9 member A7(SLC9A7) Homo sapiens This gene encodes a sodium and potassium/ proton antiporter that is a member of the solute carrier family 9 protein family. The encoded protein is primarily localized to the trans-Golgi network and is involved in maintaining pH homeostasis in organelles along the secretory and endocytic pathways. This protein may enhance cell growth of certain breast tumors. This gene is part of a gene cluster on chromosome Xp11.23. A pseudogene of this gene is found on chromosome 12. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Mar 2012],
Subcellular Localization	Golgi membrane,trans-Golgi network,plasma membrane,integral component of membrane,recycling endosome membrane,
Protein Function	function:Mediates electroneutral exchange of protons for Na(+) and K(+) across endomembranes. May contribute to Golgi volume and cation homeostasis.,miscellaneous:Is not inhibited by amiloride but by benzamil and quinine.,similarity:Belongs to the monovalent cation:proton antiporter 1 (CPA1) transporter (TC 2.A.36) family.,subunit:Interacts with SCAMP1, SCAMP2 and SCAMP5; may participate in its shuttling from trans-Golgi network to recycling endosomes.,tissue specificity:Ubiquitously expressed.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.