

Immunotag™ NMUR1 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT3159
Product Description	Immunotag™ NMUR1 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	NMUR1
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human NMUR1. AA range:1-50
Specificity	NMUR1 Polyclonal Antibody detects endogenous levels of NMUR1 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	NMUR1
Accession No.	Q9HB89 O55040
Alternate Names	NMUR1; GPR66; Neuromedin-U receptor 1; NMU-R1; G-protein coupled receptor 66; G-protein coupled receptor FM-3

Antibody Specification

Description	caution:It is uncertain whether Met-1 or Met-24 is the initiator.,function:Receptor for the neuromedin-U and neuromedin-S neuropeptides.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in greatest abundance in peripheral organs, particularly in elements of the gastrointestinal and urogenital systems with highest levels in testes. In central nervous system structures express levels are much lower than those seen in peripheral organs. Within the CNS, has been detected in highest abundance in the cerebellum, dorsal root ganglia, hippocampus, and spinal cord.,
Cell Pathway/ Category	Neuroactive ligand-receptor interaction,
Protein Expression	PNS,Testis,
Subcellular Localization	intracellular,plasma membrane,integral component of plasma membrane,integral component of membrane,
Protein Function	caution:It is uncertain whether Met-1 or Met-24 is the initiator.,function:Receptor for the neuromedin-U and neuromedin-S neuropeptides.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in greatest abundance in peripheral organs, particularly in elements of the gastrointestinal and urogenital systems with highest levels in testes. In central nervous system structures express levels are much lower than those seen in peripheral organs. Within the CNS, has been detected in highest abundance in the cerebellum, dorsal root ganglia, hippocampus, and spinal cord.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.