

Immunotag™ NRBF-2 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT3187
Product Description	Immunotag™ NRBF-2 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	NRBF-2
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	Synthesized peptide derived from NRBF-2, at AA range: 110-190
Specificity	NRBF-2 Polyclonal Antibody detects endogenous levels of NRBF-2 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	NRBF2
Accession No.	Q96F24 Q8VCQ3 Q9QYK3
Alternate Names	NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR
Description	function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand.,tissue specificity:Detected in keratinocytes, liver and placenta.,
Protein Expression	Brain,Cerebellum,Dendritic cell,Keratinocyte,Lymph,

Antibody Specification

Subcellular Localization	nucleoplasm,cytoplasm,autophagosome,cytoplasmic, membrane-bounded vesicle,cytoplasmic vesicle,phosphatidylinositol 3-kinase complex, class III,
Protein Function	function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand.,tissue specificity:Detected in keratinocytes, liver and placenta.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.