

Immunotag™ NXF3 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT3214
Product Description	Immunotag™ NXF3 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	NXF3
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human NXF3. AA range:72-121
Specificity	NXF3 Polyclonal Antibody detects endogenous levels of NXF3 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	NXF3
Accession No.	Q9H4D5
Alternate Names	NXF3; TAPL3; Nuclear RNA export factor 3; TAP-like protein 3; TAPL-3

Antibody Specification

Description	nuclear RNA export factor 3(NXF3) Homo sapiens This gene is one member of a family of nuclear RNA export factor genes. Common domain features of this family are a noncanonical RNP-type RNA-binding domain (RBD), 4 leucine-rich repeats (LRRs), a nuclear transport factor 2 (NTF2)-like domain that allows heterodimerization with NTF2-related export protein-1 (NXT1), and a ubiquitin-associated domain that mediates interactions with nucleoporins. The LRRs and NTF2-like domains are required for export activity. Alternative splicing seems to be a common mechanism in this gene family. The encoded protein of this gene has shortened LRR and ubiquitin-associated domains and its RDB is unable to bind RNA. It is located in the nucleoplasm but is not associated with either the nuclear envelope or the nucleolus. [provided by RefSeq, Jul 2008],
Protein Expression	Brain,Fetal brain,Testis,
Subcellular Localization	nucleus,nucleoplasm,cytoplasm,nuclear RNA export factor complex,
Protein Function	domain:Contains a novel CRM1-dependent nuclear export signal that compensates in cis for the loss of the nuclear pore targeting domain.,domain:Lacks C-terminal domain that mediates direct interactions with nucleoporins.,function:May function as a tissue-specific nuclear mRNA export factor.,miscellaneous:The RNA-binding domain is a non-canonical RNP-type domain.,similarity:Belongs to the NXF family.,similarity:Contains 1 LRR (leucine-rich) repeat.,similarity:Contains 1 NTF2 domain.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subcellular location:Shuttles between the nucleus and the cytoplasm.,subunit:Interacts with NXT1, NXT2, E1B-AP5 and CRM1 nuclear export factor.,tissue specificity:Expressed at high level in testis and at low level in a small number of tissues.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.