

Immunotag™ NY-CO-1 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT3215
Product Description	Immunotag™ NY-CO-1 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	NY-CO-1
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human SDCG1. AA range:881-930
Specificity	NY-CO-1 Polyclonal Antibody detects endogenous levels of NY-CO-1 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	NEMF
Accession No.	O60524 Q8CCP0
Alternate Names	NEMF; SDCCAG1; Nuclear export mediator factor NEMF; Antigen NY-CO-1; Serologically defined colon cancer antigen 1
Description	nuclear export mediator factor(NEMF) Homo sapiens This gene encodes a component of the ribosome quality control complex. The encoded protein facilitates the recognition and ubiquitination of stalled 60S subunits by the ubiquitin ligase listerin. A similar protein in fly functions as a tumor suppressor. [provided by RefSeq, Jul 2016],

Antibody Specification

Protein Expression	Kidney,Thymus,
Subcellular Localization	nucleus,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.