

## Immunotag™ OY-TES-1 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT3483
Product Description	Immunotag™ OY-TES-1 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	OY-TES-1
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,IF,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human ACRBP. AA range:181-230
Specificity	OY-TES-1 Polyclonal Antibody detects endogenous levels of OY-TES-1 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	ACRBP
Accession No.	Q8NEB7 Q3V140
Alternate Names	ACRBP; Acrosin-binding protein; Cancer/testis antigen 23; CT23; Cancer/testis antigen OY-TES-1; Proacrosin-binding protein sp32

## Antibody Specification

Description	acrosin binding protein(ACRBP) Homo sapiens The protein encoded by this gene is similar to proacrosin binding protein sp32 precursor found in mouse, guinea pig, and pig. This protein is located in the sperm acrosome and is thought to function as a binding protein to proacrosin for packaging and condensation of the acrosin zymogen in the acrosomal matrix. This protein is a member of the cancer/testis family of antigens and it is found to be immunogenic. In normal tissues, this mRNA is expressed only in testis, whereas it is detected in a range of different tumor types such as bladder, breast, lung, liver, and colon. [provided by RefSeq, Jul 2008],
Protein Expression	Testis,
Subcellular Localization	acrosomal vesicle,acrosomal membrane,extracellular region,nucleus,
Protein Function	function:May be involved in packaging and condensation of the acrosin zymogen in the acrosomal matrix via its association with proacrosin.,PTM:Phosphorylated on Tyr residues in capacitated sperm.,PTM:The N-terminus is blocked.,subcellular location:Colocalizes with proacrosin in the acrosome of sperm.,subunit:Binds proacrosin.,tissue specificity:Expression restricted to testis in normal tissue. Expressed in a wide spectrum of cancers, including bladder, breast, liver, lung and colon cancers.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.