

Immunotag™ p52 S6 kinase Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT3523
Product Description	Immunotag™ p52 S6 kinase Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	p52 S6 kinase
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human RPS6KC1. AA range:231-280
Specificity	p52 S6 kinase Polyclonal Antibody detects endogenous levels of p52 S6 kinase protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	RPS6KC1
Accession No.	Q96S38 Q8BLK9
Alternate Names	RPS6KC1; RPK118; Ribosomal protein S6 kinase delta-1; S6K-delta-1; 52 kDa ribosomal protein S6 kinase; Ribosomal S6 kinase-like protein with two PSK domains 118 kDa protein; SPHK1-binding protein

Antibody Specification

Description	<p>catalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:Instead of Lys-820, Arg-820 is found at the binding site.,domain:The first protein kinase domain appears to be a pseudokinase domain as it does not contain the classical characteristics, such as the ATP-binding motif, ATP-binding site and active site.,function:May be involved in transmitting sphingosine-1 phosphate (SPP)-mediated signaling into the cell.,similarity:Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. S6 kinase subfamily.,similarity:Contains 1 MIT domain.,similarity:Contains 1 PX (phox homology) domain.,similarity:Contains 2 protein kinase domains.,subcellular location:Also found in some small dot-like or ring-shaped early endosome structures.,subunit:Interacts with SPHK1 and phosphatidylinositol 3-phosphate.,tissue specificity:Highly expressed in testis, skeletal muscle, brain, heart, placenta, kidney and liver and weakly expressed in thymus, small intestine, lung and colon.,</p>
Cell Pathway/ Category	Regulates Angiogenesis, Insulin Receptor, B Cell Receptor, AMPK
Protein Expression	Amygdala,Brain,Epithelium,
Subcellular Localization	cytoplasm,early endosome,ribosome,membrane,intracellular membrane-bounded organelle,
Protein Function	<p>catalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:Instead of Lys-820, Arg-820 is found at the binding site.,domain:The first protein kinase domain appears to be a pseudokinase domain as it does not contain the classical characteristics, such as the ATP-binding motif, ATP-binding site and active site.,function:May be involved in transmitting sphingosine-1 phosphate (SPP)-mediated signaling into the cell.,similarity:Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. S6 kinase subfamily.,similarity:Contains 1 MIT domain.,similarity:Contains 1 PX (phox homology) domain.,similarity:Contains 2 protein kinase domains.,subcellular location:Also found in some small dot-like or ring-shaped early endosome structures.,subunit:Interacts with SPHK1 and phosphatidylinositol 3-phosphate.,tissue specificity:Highly expressed in testis, skeletal muscle, brain, heart, placenta, kidney and liver and weakly expressed in thymus, small intestine, lung and colon.,</p>
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.