

Immunotag™ PP4R2 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT3834
Product Description	Immunotag™ PP4R2 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PP4R2
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Monkey
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human PPP4R2. AA range:171-220
Specificity	PP4R2 Polyclonal Antibody detects endogenous levels of PP4R2 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	PPP4R2
Accession No.	Q9NY27 Q0VGB7
Alternate Names	PPP4R2; SBBI57; Serine/threonine-protein phosphatase 4 regulatory subunit 2

Antibody Specification

Description	protein phosphatase 4 regulatory subunit 2(PPP4R2) Homo sapiens The protein encoded by this gene is a regulatory subunit of the serine/threonine-protein phosphatase 4 complex. In addition to being required for efficient DNA double strand break repair, this complex plays a role in organization of microtubules at centrosomes and processing of spliceosomal snRNPs. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015],
Protein Expression	Corpus callosum,Uterus,
Subcellular Localization	nucleus,nucleoplasm,cytoplasm,centrosome,protein phosphatase 4 complex,
Protein Function	function:Regulatory subunit of serine/threonine-protein phosphatase 4 (PP4). May regulate the activity of PPP4C at centrosomal microtubule organizing centers. Its interaction with the SMN complex leads to enhance the temporal localization of snRNPs, suggesting a role of PPP4C in maturation of spliceosomal snRNPs. The PPP4C-PPP4R2-PPP4R3A PP4 complex specifically dephosphorylates H2AFX phosphorylated on 'Ser-140' (gamma-H2AFX) generated during DNA replication and required for DNA DSB repair.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Belongs to the PPP4R2 family.,subunit:Serine/threonine-protein phosphatase 4 (PP4) occurs in different assemblies of the catalytic and one or more regulatory subunits. Component of the PP4 complexes PPP4C-PPP4R2, PPP4C-PPP4R2-PPP4R3A and PPP4C-PPP4R2-PPP4R3B. The PPP4C-PPP4R2 complex appears to be a tetramer composed of 2 molecules of PPP4C and 2 molecules of PPP4R2. Interacts with DDX20/GEMIN3 and GEMIN4.,tissue specificity:Widely expressed.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.