

Immunotag™ PXR Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT3911
Product Description	Immunotag™ PXR Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PXR
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human NR1I2. AA range:91-140
Specificity	PXR Polyclonal Antibody detects endogenous levels of PXR protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	NR1I2
Accession No.	O75469 O54915
Alternate Names	NR1I2; PXR; Nuclear receptor subfamily 1 group I member 2; Orphan nuclear receptor PAR1; Orphan nuclear receptor PXR; Pregnane X receptor; Steroid and xenobiotic receptor; SXR

Antibody Specification

Description	nuclear receptor subfamily 1 group I member 2(NR1I2) Homo sapiens This gene product belongs to the nuclear receptor superfamily, members of which are transcription factors characterized by a ligand-binding domain and a DNA-binding domain. The encoded protein is a transcriptional regulator of the cytochrome P450 gene CYP3A4, binding to the response element of the CYP3A4 promoter as a heterodimer with the 9-cis retinoic acid receptor RXR. It is activated by a range of compounds that induce CYP3A4, including dexamethasone and rifampicin. Several alternatively spliced transcripts encoding different isoforms, some of which use non-AUG (CUG) translation initiation codon, have been described for this gene. Additional transcript variants exist, however, they have not been fully characterized. [provided by RefSeq, Jul 2008],
Protein Expression	Hepatoma,Liver,Peripheral blood,
Subcellular Localization	nucleus,nucleoplasm,
Protein Function	function:Orphan receptor; its natural ligand is probably pregnane. Binds to a response element in the CYP3A4 and ABCB1/MDR1 genes promoter. Activates its expression in response to a wide variety of endobiotics and xenobiotics.,induction:Activated by naturally occurring steroids such as pregnenolone and progesterone.,similarity:Belongs to the nuclear hormone receptor family. NR1 subfamily.,similarity:Contains 1 nuclear receptor DNA-binding domain.,subunit:Forms a heterodimer with RXR.,tissue specificity:Expressed in liver, colon and small intestine.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.