

Immunotag™ Rabenosyn-5 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT3949
Product Description	Immunotag™ Rabenosyn-5 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Rabenosyn-5
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human ZFYVE20. AA range:145-194
Specificity	Rabenosyn-5 Polyclonal Antibody detects endogenous levels of Rabenosyn-5 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	ZFYVE20
Accession No.	Q9H1K0 Q80Y56
Alternate Names	ZFYVE20; Rabenosyn-5; 110 kDa protein; FYVE finger-containing Rab5 effector protein rabenosyn-5; Zinc finger FYVE domain-containing protein 20

Antibody Specification

Description	rabenosyn, RAB effector(RBSN) Homo sapiens This gene encodes a protein that belongs to the FYVE zinc finger family of proteins. The encoded protein interacts with Ras-related proteins that regulate membrane trafficking. A missense mutation in this gene is associated with a defect in the early endocytic pathway. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2014],
Cell Pathway/ Category	Endocytosis,
Protein Expression	Brain,Cervix carcinoma,Epithelium,Human salivary gland,
Subcellular Localization	endosome,cytosol,plasma membrane,endosome membrane,cytoplasmic side of endosome membrane,early endosome membrane,intracellular membrane-bounded organelle,extracellular exosome,
Protein Function	function:Rab4/Rab5 effector protein acting in early endocytic membrane fusion and membrane trafficking of recycling endosomes. Required for endosome fusion either homotypically or with clathrin coated vesicles. Plays a role in the lysosomal trafficking of cathepsin D from the Golgi to lysosomes. Also promotes the recycling of transferrin directly from early endosomes to the plasma membrane. Binds phospholipid vesicles containing phosphatidylinositol 3-phosphate (PtdInsP3).,similarity:Contains 1 C2H2-type zinc finger.,similarity:Contains 1 FYVE-type zinc finger.,similarity:Contains 1 UIM (ubiquitin-interacting motif) repeat.,subunit:Interacts with EHD1, RAB4A, RAB5A, RAB22A, RAB24 and VPS45. Binds simultaneously to RAB4A and RAB5A in vitro. Interacts with RAB4A and RAB5A that has been activated by GTP binding.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.