

Immunotag™ Reg IV Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT4040
Product Description	Immunotag™ Reg IV Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Reg IV
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	IHC-p,ELISA
Recommended Dilution	Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	Synthesized peptide derived from the Internal region of human Reg IV.
Specificity	Reg IV Polyclonal Antibody detects endogenous levels of Reg IV protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	REG4
Accession No.	Q9BYZ8 Q9D8G5
Alternate Names	REG4; GISP; RELP; Regenerating islet-derived protein 4; REG-4; Gastrointestinal secretory protein; REG-like protein; Regenerating islet-derived protein IV; Reg IV

Antibody Specification

Description	function:May be involved in inflammatory and metaplastic responses of the gastrointestinal epithelium.,induction:Up-regulated by mucosal injury from active Crohn's disease or ulcerative colitis. Up-regulated in colorectal tumors. Up-regulated in epithelial cells at regenerating margins of peptic ulcers in the stomach and duodenum.,online information:Regenerating protein IV,similarity:Contains 1 C-type lectin domain.,tissue specificity:Highly expressed in the gastrointestinal tract including the duodenum, jejunum, ileum, ileocecum, appendix, descending colon, pancreas and small intestine. Weakly expressed in normal colon and stomach. Strongly expressed in most colorectal tumors than in normal colon. Preferentially expressed in mucinous tumors and in some cases neuro-endocrine tumors. Expressed in mucus-secreting cells and enterocyte-like cells. In small intestine expressed at the basal perinuclear zone of goblet cells.,
Protein Expression	Colon,Colon cancer,
Subcellular Localization	extracellular region,cytoplasm,
Protein Function	function:May be involved in inflammatory and metaplastic responses of the gastrointestinal epithelium.,induction:Up-regulated by mucosal injury from active Crohn's disease or ulcerative colitis. Up-regulated in colorectal tumors. Up-regulated in epithelial cells at regenerating margins of peptic ulcers in the stomach and duodenum.,online information:Regenerating protein IV,similarity:Contains 1 C-type lectin domain.,tissue specificity:Highly expressed in the gastrointestinal tract including the duodenum, jejunum, ileum, ileocecum, appendix, descending colon, pancreas and small intestine. Weakly expressed in normal colon and stomach. Strongly expressed in most colorectal tumors than in normal colon. Preferentially expressed in mucinous tumors and in some cases neuro-endocrine tumors. Expressed in mucus-secreting cells and enterocyte-like cells. In small intestine expressed at the basal perinuclear zone of goblet cells.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.