

# Immunotag™ Relaxin 3 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT4041
Product Description	Immunotag™ Relaxin 3 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Relaxin 3
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	IHC-p,ELISA
Recommended Dilution	Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	Synthesized peptide derived from the Internal region of human Relaxin 3.
Specificity	Relaxin 3 Polyclonal Antibody detects endogenous levels of Relaxin 3 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	RLN3
Accession No.	Q8WXF3 Q8CHK2
Alternate Names	RLN3; INSL7; RXN3; ZINS4; Relaxin-3; Insulin-like peptide INSL7; Insulin-like peptide 7; Prorelaxin H3

## Antibody Specification

Description	relaxin 3(RLN3) Homo sapiens This gene encodes a member of the relaxin family of insulin-like hormones that is expressed predominantly in the brain and plays a role in physiological processes such as stress, memory and appetite regulation. The encoded protein is a precursor that is proteolytically processed to generate a heterodimeric mature form consisting A and B chains interlinked by disulfide bonds. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2015],
Subcellular Localization	extracellular region,
Protein Function	function:May play a role in neuropeptide signaling processes. Ligand for LGR7, relaxin-3 receptor-1 (GPCR135) and relaxin-3 receptor-2 (GPCR142).,similarity:Belongs to the insulin family.,subunit:Heterodimer of a B chain and an A chain linked by two disulfide bonds.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.