

Immunotag™ SLK Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT4320
Product Description	Immunotag™ SLK Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	SLK
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,IF,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	Synthesized peptide derived from SLK, at AA range: 1120-1200
Specificity	SLK Polyclonal Antibody detects endogenous levels of SLK protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	SLK
Accession No.	Q9H2G2 O54988 O08815
Alternate Names	SLK; KIAA0204; STK2; STE20-like serine/threonine-protein kinase; STE20-like kinase; hSLK; CTCL tumor antigen se20-9; STE20-related serine/threonine-protein kinase; STE20-related kinase; Serine/threonine-protein kinase 2

Antibody Specification

Description	<p>catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Mediates apoptosis and actin stress fiber dissolution.,PTM:Autophosphorylated. Phosphorylated upon DNA damage, probably by ATM or ATR.,PTM:Proteolytically cleaved by caspase-3.,sequence caution:Contaminating sequence. Potential poly-A sequence starting in position 422.,sequence caution:Contaminating sequence. Potential poly-A sequence starting in position 611.,sequence caution:Contaminating sequence. Potential poly-A sequence starting in position 614.,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. STE20 subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 UVR domain.,tissue specificity:Ubiquitously expressed. Highest expression is found in heart and in skeletal muscle.,</p>
Cell Pathway/ Category	Oocyte meiosis,
Protein Expression	Bone marrow,Epithelium,Lung carcinoma,Testis,Uterus,
Subcellular Localization	cytoplasm,cell-cell adherens junction,cell leading edge,perinuclear region of cytoplasm,extracellular exosome,
Protein Function	<p>catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Mediates apoptosis and actin stress fiber dissolution.,PTM:Autophosphorylated. Phosphorylated upon DNA damage, probably by ATM or ATR.,PTM:Proteolytically cleaved by caspase-3.,sequence caution:Contaminating sequence. Potential poly-A sequence starting in position 422.,sequence caution:Contaminating sequence. Potential poly-A sequence starting in position 611.,sequence caution:Contaminating sequence. Potential poly-A sequence starting in position 614.,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. STE20 subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 UVR domain.,tissue specificity:Ubiquitously expressed. Highest expression is found in heart and in skeletal muscle.,</p>
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.