

Immunotag™ StARD13 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT4438
Product Description	Immunotag™ StARD13 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	StARD13
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human STA13. AA range:101-150
Specificity	StARD13 Polyclonal Antibody detects endogenous levels of StARD13 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	STARD13
Accession No.	Q9Y3M8 Q923Q2
Alternate Names	STARD13; DLC2; GT650; StAR-related lipid transfer protein 13; 46H23.2; Deleted in liver cancer 2 protein; DLC-2; Rho GTPase-activating protein; START domain-containing protein 13; StARD13

Antibody Specification

Description	StAR related lipid transfer domain containing 13(STARD13) Homo sapiens This gene encodes a protein which contains an N-terminal sterile alpha motif (SAM) for protein-protein interactions, followed by an ATP/GTP-binding motif, a GTPase-activating protein (GAP) domain, and a C-terminal STAR-related lipid transfer (START) domain. It may be involved in regulation of cytoskeletal reorganization, cell proliferation, and cell motility, and acts as a tumor suppressor in hepatoma cells. The gene is located in a region of chromosome 13 that is associated with loss of heterozygosity in hepatocellular carcinomas. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Aug 2011],
Protein Expression	Human salivary gland,Skin,Uterus,
Subcellular Localization	lipid particle,cytosol,mitochondrial membrane,
Protein Function	function:GTPase-activating protein for RhoA, and perhaps for Cdc42. May be involved in regulation of cytoskeletal reorganization, cell proliferation and cell motility. Acts a tumor suppressor in hepatocellular carcinoma cells.,similarity:Contains 1 Rho-GAP domain.,similarity:Contains 1 SAM (sterile alpha motif) domain.,similarity:Contains 1 START domain.,subunit:Homodimer. Interacts with TAX1BP1.,tissue specificity:Ubiquitously expressed. Underexpressed in hepatocellular carcinoma cells and some breast cancer cell lines.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.