

Immunotag™ Synaptotagmin XIII Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT4488
Product Description	Immunotag™ Synaptotagmin XIII Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Synaptotagmin XIII
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	Synthesized peptide derived from Synaptotagmin XIII, at AA range: 290-370
Specificity	Synaptotagmin XIII Polyclonal Antibody detects endogenous levels of Synaptotagmin XIII protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	SYT13
Accession No.	Q7L8C5 Q9EQT6 Q925B5
Alternate Names	SYT13; KIAA1427; Synaptotagmin-13; Synaptotagmin XIII; SytXIII

Antibody Specification

Description	synaptotagmin 13(SYT13) Homo sapiens This gene encodes a member of the large synaptotagmin protein family. Family members have an extracellular N-terminal transmembrane domain and a cytoplasmic C terminus with two tandem C2 domains (C2A and C2B). Synaptotogmin family members can form homo- and heteromeric complexes with each other. They also have different biochemical properties and developmental profiles, and patterns of tissue distribution. Synaptotagmins function as membrane traffickers in multicellular organisms. Two alternatively spliced transcript variants that encode different protein isoforms have been described for this gene. [provided by RefSeq, Oct 2011],
Protein Expression	Brain,Fetal brain,
Subcellular Localization	plasma membrane,integral component of plasma membrane,transport vesicle,intracellular membrane-bounded organelle,presynapse,
Protein Function	domain:The first C2 domain/C2A does not mediate Ca(2+)-dependent phospholipid binding.,domain:The second C2 domain/C2B domain binds phospholipids regardless of whether calcium is present.,function:May be involved in transport vesicle docking to the plasma membrane.,similarity:Belongs to the synaptotagmin family.,similarity:Contains 2 C2 domains.,subunit:Interacts with NRXN1.,tissue specificity:Expressed in brain, pancreas and kidney.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.