

Immunotag™ T2R16 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT4507
Product Description	Immunotag™ T2R16 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	T2R16
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human TAS2R16. AA range:136-185
Specificity	T2R16 Polyclonal Antibody detects endogenous levels of T2R16 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	TAS2R16
Accession No.	Q9NYV7 P59529
Alternate Names	TAS2R16; Taste receptor type 2 member 16; T2R16

Antibody Specification

Description	taste 2 receptor member 16(TAS2R16) Homo sapiens This gene encodes a member of a family of candidate taste receptors that are members of the G protein-coupled receptor superfamily. These family members are specifically expressed by taste receptor cells of the tongue and palate epithelia. Each of these apparently intronless genes encodes a 7-transmembrane receptor protein, functioning as a bitter taste receptor. This gene is clustered with another 3 candidate taste receptor genes in chromosome 7 and is genetically linked to loci that influence bitter perception. [provided by RefSeq, Jul 2008],
Cell Pathway/ Category	Taste transduction,
Subcellular Localization	endoplasmic reticulum,trans-Golgi network,plasma membrane,external side of plasma membrane,integral component of membrane,
Protein Function	function:Gustducin-coupled receptor implicated in the perception of bitter compounds in the oral cavity and the gastrointestinal tract. Signals through PLCB2 and the calcium-regulated cation channel TRPM5.,miscellaneous:Confers bitter perception of salicin to non-taster mice.,miscellaneous:Several bitter taste receptors are expressed in a single taste receptor cell.,polymorphism:The Lys-172 polymorphism in TAS2R16 is associated with genetic susceptibility to alcoholism [MIM:103780].,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in a subset of gustducin-positive taste receptor cells of the tongue.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.