

Immunotag™ TAF II p135/p105 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT4527
Product Description	Immunotag™ TAF II p135/p105 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	TAF II p135/p105
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human TAF4. AA range:941-990
Specificity	TAF II p135/p105 Polyclonal Antibody detects endogenous levels of TAF II p135/p105 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	TAF4/TAF4B
Accession No.	O00268/Q92750
Alternate Names	TAF4; TAF2C; TAF2C1; TAF4A; TAFII130; TAFII135; Transcription initiation factor TFIID subunit 4; RNA polymerase II TBP-associated factor subunit C; TBP-associated factor 4; Transcription initiation factor TFIID 130 kDa subunit; TAF(II)130;

Antibody Specification

Description	TATA-box binding protein associated factor 4(TAF4) Homo sapiens Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes one of the larger subunits of TFIID that has been shown to potentiate transcriptional activation by retinoic acid, thyroid hormone and
Cell Pathway/ Category	Basal transcription factors,Huntington's disease,
Subcellular Localization	nuclear chromatin,nucleoplasm,transcription factor TFIID complex,cytoplasm,transcription factor TFTC complex,MLL1 complex,
Protein Function	function:Makes part of TFIID is a multimeric protein complex that plays a central role in mediating promoter responses to various activators and repressors. Potentiates transcriptional activation by the AF-2S of the retinoic acid, vitamin D3 and thyroid hormone.,similarity:Belongs to the TAF4 family.,similarity:Contains 1 TAFH (NHR1) domain.,subunit:TFIID is composed of TATA binding protein (TBP) and a number of TBP-associated factors (TAFs). Component of the TFTC-HAT complex, at least composed of TAF5L, TAF6L, TADA3L, SUPT3H, TAF2, TAF4, TAF5, GCN5L2/GCN5, TAF10 and TRRAP. Interacts with SV40 Large T antigen.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.