

# Immunotag™ TAF II p18 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT4528
Product Description	Immunotag™ TAF II p18 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	TAF II p18
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	IHC-p,ELISA
Recommended Dilution	Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human TAF13. AA range:71-120
Specificity	TAF II p18 Polyclonal Antibody detects endogenous levels of TAF II p18 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	TAF13
Accession No.	Q15543 P61216
Alternate Names	TAF13; TAF2K; TAFII18; Transcription initiation factor TFIID subunit 13; Transcription initiation factor TFIID 18 kDa subunit; TAF(II)18; TAFII-18; TAFII18

## Antibody Specification

Description	TATA-box binding protein associated factor 13(TAF13) Homo sapiens Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes a small subunit associated with a subset of TFIID complexes. This subunit interacts with TBP and with two other small subunits of TFIID
Cell Pathway/ Category	Basal transcription factors,
Protein Expression	PCR rescued clones,
Subcellular Localization	nucleus,nucleoplasm,transcription factor TFIID complex,nucleolus,
Protein Function	domain:The binding of TAF10 and TAF11 requires distinct domains of TAF13.,function:TFIID beta-specific TAFII.,similarity:Belongs to the TAF13 family.,similarity:Contains 1 histone-fold domain.,subunit:TFIID is composed of TATA binding protein (TBP) and a number of TBP-associated factors (TAFs). Interacts with TBP, and more strongly with TAF10 and TAF11.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.