

Immunotag™ TF Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT4610
Product Description	Immunotag™ TF Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	TF
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Monkey
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human Coagulation Factor III. AA range:246-295
Specificity	TF Polyclonal Antibody detects endogenous levels of TF protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	F3
Accession No.	P13726 P20352
Alternate Names	F3; Tissue factor; TF; Coagulation factor III; Thromboplastin; CD antigen CD142

Antibody Specification

Description	coagulation factor III, tissue factor(F3) Homo sapiens This gene encodes coagulation factor III which is a cell surface glycoprotein. This factor enables cells to initiate the blood coagulation cascades, and it functions as the high-affinity receptor for the coagulation factor VII. The resulting complex provides a catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades, which circulate as nonfunctional precursors, this factor is a potent initiator that is fully functional when expressed on cell surfaces. There are 3 distinct domains of this factor: extracellular, transmembrane, and cytoplasmic. This protein is the only one in the coagulation pathway for which a congenital deficiency has not been described. Alternate splicing results in multiple transcript variants.[provided by RefSeq, May 2010],
Cell Pathway/ Category	Complement and coagulation cascades,
Protein Expression	Lung,Prostate,
Subcellular Localization	extracellular space,plasma membrane,cell surface,integral component of membrane,extracellular matrix,intrinsic component of external side of plasma membrane,extracellular exosome,serine-type peptidase complex,
Protein Function	function:Initiates blood coagulation by forming a complex with circulating factor VII or VIIa. The [TF:VIIa] complex activates factors IX or X by specific limited proteolysis. TF plays a role in normal hemostasis by initiating the cell-surface assembly and propagation of the coagulation protease cascade.,induction:TF expression is highly dependent upon cell type. TF can also be induced by the inflammatory mediators interleukin 1 and TNF, as well as by endotoxin, to appear on monocytes and vascular endothelial cells as a component of cellular immune response.,online information:The Singapore human mutation and polymorphism database,online information:Tissue factor entry,similarity:Belongs to the tissue factor family.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.